

Neural Network-Based Methods for Inverse Problems in Holography

Hyun-Sik Jeong

APCTP

Sogang University, 29 December 2025

The talk is mainly a **review**, but here are some of my contributions

Holographic CMT, QCD

arXiv:2401.00939 [pdf, other] hep-th cond-mat.dis-nn gr-qc hep-ph doi 10.1007/JHEP03(2024)141

Deep learning bulk spacetime from boundary optical conductivity

Authors: Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

arXiv:2502.10245 [pdf, other] hep-th cond-mat.dis-nn cond-mat.str-el gr-qc doi 10.1103/nh4n-ylnz (PRD)

Deep learning-based holography for T-linear resistivity

Authors: Byoungjoon Ahn, Hyun-Sik Jeong, Chang-Woo Ji, Keun-Young Kim, Kwan Yun

arXiv:2511.22522 [pdf, ps, other] hep-th cs.LG

AdS/Deep-Learning made easy II: neural network-based approaches to holography and inverse problems


Authors: Hyun-Sik Jeong, Hanse Kim, Keun-Young Kim, Gaya Yun, Hyeonwoo Yu, Kwan Yun

Holographic QI


arXiv:2406.07395 [pdf, other] hep-th cond-mat.dis-nn gr-qc doi 10.1007/JHEP01(2025)025

Holographic reconstruction of black hole spacetime: machine learning and entanglement entropy

Authors: Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun



Quantum Information Black Holes & Holography



“Quantum Gravity & Quantum Many-Body”

PostDoc



Ashutosh Tripathi

PostDoc




Yongjun Ahn

PostDoc



???



Sep 2025 - Present

apctp

JRG Team




Today's Talk

(Inverse Problems in Holography)


Holography

Holographic Duality

Strongly
Coupled



Perturbation
Theory (X)



Ex: AdS/CFT Correspondence

- 1997 – [Juan Maldacena](#) proposed the [AdS/CFT correspondence](#)

Top Cited Articles of All Time

As of 2025

The Large N limit of superconformal field theories and supergravity

AdS/CFT

#1

Juan Martin Maldacena (Harvard U.) (Nov, 1997)

Published in: *Int.J.Theor.Phys.* 38 (1999) 1113-1133 (reprint), *Adv.Theor.Math.Phys.* 2 (1998) 231-252 • e-Print: [hep-th/9711200](#) [hep-th]

[pdf](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[↻](#) 20,774 citations

A Model of Leptons

#2

Steven Weinberg (MIT, LNS) (Nov, 1967)

Published in: *Phys.Rev.Lett.* 19 (1967) 1264-1266

[links](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[↻](#) 14,900 citations

Anti de Sitter space and holography

AdS/CFT

#3

Edward Witten (Princeton, Inst. Advanced Study) (Feb, 1998)

Published in: *Adv.Theor.Math.Phys.* 2 (1998) 253-291 • e-Print: [hep-th/9802150](#) [hep-th]

[pdf](#) [links](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[↻](#) 13,246 citations

Gauge theory correlators from noncritical string theory

#4

S.S. Gubser (Princeton U.), Igor R. Klebanov (Princeton U.), Alexander M. Polyakov (Princeton U.) (Feb, 1998)

Published in: *Phys.Lett.B* 428 (1998) 105-114 • e-Print: [hep-th/9802109](#) [hep-th]

[pdf](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[↻](#) 10,909 citations

CP Conservation in the Presence of Instantons

#5

R.D. Peccei (Stanford U., ITP), Helen R. Quinn (Stanford U., ITP) (Mar, 1977)

Published in: *Phys.Rev.Lett.* 38 (1977) 1440-1443

[links](#) [DOI](#) [cite](#) [claim](#)

[reference search](#)

[↻](#) 8,540 citations

Juan Maldacena



Born 10 September 1968 (age 56)
Buenos Aires, Argentina

Education [Balseiro Institute \(BS\)](#)
[Princeton University \(PhD\)](#)

Known for [AdS/CFT correspondence](#)
[ER = EPR](#)
[ABJM superconformal field theory](#)

Awards [Alfred P. Sloan Foundation Fellowship](#)
[Sackler Prize](#)
[MacArthur Fellowship](#)
[Xanthopoulos Pius XI Medal](#)
[Dirac Medal](#)

No Proof Yet

Evidence of AdS/CFT


String Theory, SUSY

- SYM theory in (3+1) dimensions $\longleftrightarrow$ IIB supergravity on $\text{AdS}_5 \times S^5$
- ABJM theory in (2+1) dimensions $\longleftrightarrow$ 11d supergravity on $\text{AdS}_4 \times S^7/\mathbb{Z}_k$
- SCFT theory in (1+1) dimensions $\longleftrightarrow$ IIB supergravity on $\text{AdS}_3 \times S^3 \times M^4$

Entanglement Entropy


- Explicit example of AdS/CFT
- Measure of entanglement
- Quantum nature of gravity

Nobel Prize in Physics 2022



© Johan Jarnestad/The Royal Swedish Academy of Sciences

Ryu-Takayanagi Formula





Holographic Derivation of Entanglement Entropy from the anti-de Sitter Space/Conformal Field Theory Correspondence


Shinsei Ryu and Tadashi Takayanagi
Phys. Rev. Lett. **96**, 181602 – Published 9 May 2006

Physical Review Letters

Renyi entropy
Reflected entropy
Krylov complexity
...



Shinsei Ryu



Tadashi Takayanagi

$$S_A = \frac{\text{Area}(\gamma_A)}{4G_N}$$


Quantum Gravity

(Quantum) entanglement entropy
= Area of extremal surface

“New perspective of the nature of gravity”


Applied Holography

Gravity Theory




Quantum Physics

Describing gravitational systems
(e.g., Black Holes)




Describing **strongly interacting**
quantum systems




Perturbation Theory (X)

- Study **quantum gravity** in AdS (fundamental physics)
- Study **strongly interacting physics** (practical physics)

Holographic Methods for Many-Body Physics



Holographists use a string theory correspondence that relates black holes and condensed-matter systems, such as metals and superconductors.



Strongly Interacting
Quark-Gluon Plasma



High-Tc Superconductors

- Holography has generated a **new insights** and **perspective** on many-body physics problem
- Holography has prompted efforts from **field theory**, **hydrodynamics** and even **experiments**

e.g, **anomalous transport phenomena**

Stranger than metals REVIEW

PHILIP W. PHILLIPS , NIGEL E. HUSSEY , AND PETER ABBAMONTE  [Authors Info & Affiliations](#)

SCIENCE • 8 Jul 2022 • Vol 377, Issue 6602 • DOI: 10.1126/science.abh4273






Table 2. Snapshot of current theoretical modeling of the strange metal regime. Indicated are consistency with T -linear resistivity, $\omega^{-2/3}$ scaling of the mid-IR optical conductivity, quadrature-scaling magnetoresistance, extended quantum criticality, and what predictions are made in terms of experimental observables. Scenarios: MFL, marginal Fermi liquid; EFL, ersatz Fermi liquid; SYK, Sachdev-Ye-Kitaev; AdS/CFT, anti-de Sitter space/conformal field theory conjecture; AD/EMD, anomalous dimensions/Einstein-Maxwell-dilaton; HM, Hubbard model; QMC, quantum Monte Carlo;

ED, exact diagonalization; CA, cold atoms; DMFT/EDMFT, dynamical mean-field theory/embedded dynamical mean-field theory; A-B, Aharonov-Bohm effect; ECFL, extremely correlated Fermi liquid; QCP, quantum critical point. * T -linear resistivity is an input. **A slope change occurs through the MIR. ***Quadrature scaling obtained only for a bivalued random resistor model (121) with equal weights (27). ****Although this scaling was thought to arise in pure AdS with an inhomogeneous charge density (123), later studies (124, 125) found otherwise.

T-linear Resistivity

$$\rho \propto T \text{ as } T \rightarrow 0$$

$$\rho \propto T \text{ as } T \rightarrow \infty$$

$$\sigma \propto \omega^{-2/3}$$

Quadrature MR

Extended criticality


Experimental prediction

	$\rho \propto T \text{ as } T \rightarrow 0$	$\rho \propto T \text{ as } T \rightarrow \infty$	$\sigma \propto \omega^{-2/3}$	Quadrature MR	Extended criticality	Experimental prediction
<i>Phenomenological</i>						
MFL	$\sqrt{\text{ (67)}}$	$\times \text{ (67)}$	$\times$	$\times$	$\times$	Loop currents (107)
EFL	— *	—	—	$\times$	$\times$	Loop currents (108)
<i>Numerical</i>						
ECFL	$\times$	(109)	—	—	$\times$	$\times$
HM (QMC/ED/CA)	— (110)	$\sqrt{\text{ (110–114)}}$	$\times$	—	—	—
DMFT/EDMFT	$\sqrt{\text{ (115)}}$	$\sqrt{\text{ (116, 117)}}$	$\times$	—	$\sqrt{\text{ (117)}}$	—
QCP	(118)	—	—	—	$\times$	—
<i>Gravity-based</i>						
SYK	$\sqrt{\text{ (119, 120)}}$	$\sqrt{**} \text{ (120)}$	$\times$	$\sqrt{****} \text{ (121)}$	—	$\times$
AdS/CFT	$\sqrt{\text{ (122)}}$	$\sqrt{\text{ (122)}}$	$\sqrt{*****} \text{ (90, 126)}$	$\times$	$\times$	$\times$
AD/EMD	$\sqrt{\text{ (127–129)}}$	$\sqrt{\text{ (90, 126, 127, 129, 130)}}$	$\sqrt{\text{ (90, 126, 130)}}$	$\times$	$\sqrt{\text{ (126)}}$	Fractional A-B (129)

Holography



EX 1: Quark-Gluon Plasma



PHYSICAL REVIEW LETTERS

Highlights Recent Accepted Collections Authors Referees Search Press About Editorial Team

Viscosity in Strongly Interacting Quantum Field Theories from Black Hole Physics

P. K. Kovtun, D. T. Son, and A. O. Starinets



Phys. Rev. Lett. **94**, 111601 – Published 22 March 2005

- Viscosity of Quark-Gluon Plasma (Insights into the early universe / heavy-ion collisions)


Perturbative Theory: $\eta/s = \frac{A}{\lambda^2 \log(B/\sqrt{\lambda})} \gg 1$ (small 't Hooft coupling λ)

AdS/QCD Theory: $\eta/s = 1/(4\pi)$

Experiment: $\eta/s \approx 0.19$



EX 2: Superconductors



PHYSICAL REVIEW LETTERS

Highlights Recent Accepted Collections Authors Referees Search Press About Editorial Team

Building a Holographic Superconductor

Sean A. Hartnoll, Christopher P. Herzog, and Gary T. Horowitz
Phys. Rev. Lett. **101**, 031601 – Published 14 July 2008

- New perspective of high- T_c superconductivity
- Spontaneous condensation, infinite conductivities, ...

$$\sigma(\omega) = \sigma_0 + \left(\frac{i}{\omega} + \delta(\omega) \right) \frac{\rho_s}{\mu}$$



- Energy gap

BCS Theory: $\omega_g/T_c \approx 3.5$ (Weakly-coupled BCS Theory)

AdS/CMT Theory: $\omega_g/T_c \approx 8$ (Holographic Superconductors)

Experiment: $\omega_c/T_c \approx 7.9 \pm 0.5$ (High- T_c Cuprate Superconductors)



[Nature](#) **447**, 569–572 (2007)



AdS/Deep Learning


Forward Holography

(Standard Method)




Data (fitted by some model)

Inverse Holography



Scientific Machine Learning (SciML)

Machine Learning



Scientific Knowledge



Scientific Machine Learning

Data-driven methods

More powerful, robust, interpretable ML modeling

Terminology

Scientific Machine Learning

Sub-field

vs. purely data-driven approach

Physics-Informed Machine Learning

(+ **Physics laws & equations** into the learning process)

PDE / ODE

**Physics-Informed
Neural Networks**

PINNs

ODE

**Neural Ordinary
Differential Equations**

Neural ODEs

Useful References

nature reviews physics

Review Article | Published: 24 May 2021

Physics-informed machine learning

[George Em Karniadakis](#) , [Ioannis G. Kevrekidis](#), [Lu Lu](#), [Paris Perdikaris](#), [Sifan Wang](#) & [Liu Yang](#)

[Nature Reviews Physics](#) **3**, 422–440 (2021) | [Cite this article](#)

PIML

REVIEWS OF MODERN PHYSICS

Machine learning and the physical sciences^{*}

Giuseppe Carleo, Ignacio Cirac, Kyle Cranmer, Laurent Daudet, Maria Schuld, Naftali Tishby, Leslie Vogt-Maranto, and Lenka Zdeborová

Rev. Mod. Phys. **91**, 045002 – Published 6 December 2019

Physics

Science

Solving the quantum many-body problem with artificial neural networks

[GIUSEPPE CARLEO](#)  AND [MATTHIAS TROYER](#) [Authors Info & Affiliations](#)

SCIENCE • 10 Feb 2017 • Vol 355, Issue 6325 • pp. 602-606 • DOI: 10.1126/science.aag2302

Many-Body Physics

TOPICAL REVIEW

Machine learning for condensed matter physics

Edwin Bedolla¹ , Luis Carlos Padierna^{2,1}  and Ramón Castañeda-Priego¹ 

Published 5 November 2020 • © 2020 IOP Publishing Ltd

[Journal of Physics: Condensed Matter](#), [Volume 33](#), [Number 5](#)

Physics Reports

Data science applications to string theory

[Fabian Ruehle](#)  

[Volume 839](#), 21 January 2020

String Theory

Broad Applicability of SciML

Particle physics

Many-body physics


Cosmology

Quantum computing

String theory

...

My Interests..



AdS/DL Correspondence

Physical Review D

OPEN ACCESS | GO MOBILE » | ACCESS BY POSTECH

Deep learning and the AdS / CFT correspondence

[Koji Hashimoto](#)¹, [Sotaro Sugishita](#)¹, [Akinori Tanaka](#)^{2,3,4}, and [Akio Tomiya](#)⁵

Phys. Rev. D **98**, 046019 – Published 27 August, 2018

DOI: <https://doi.org/10.1103/PhysRevD.98.046019>

Abstract

We present a deep neural network representation of the AdS / CFT correspondence, and demonstrate the emergence of the bulk metric function via the learning process for given data sets of response in boundary quantum field theories. The emergent radial direction of the bulk is identified with the depth of the layers, and the network itself is interpreted as a bulk geometry. Our network provides a data-driven holographic modeling of strongly coupled systems. With a scalar ϕ^4 theory with unknown mass and coupling, in unknown curved spacetime with a black hole horizon, we demonstrate that our deep learning (DL) framework can determine the systems that fit given response data. First, we show that, from boundary data generated by the anti-de Sitter (AdS) Schwarzschild spacetime, our network can reproduce the metric. Second, we demonstrate that our network with experimental data as an input can determine the bulk metric, the mass and the quadratic coupling of the holographic model. As an example we use the experimental data of the magnetic response of the strongly correlated material $\text{Sm}_{0.6}\text{Sr}_{0.4}\text{MnO}_3$. This AdS / DL correspondence not only enables gravitational modeling of strongly correlated systems, but also sheds light on a hidden mechanism of the emerging space in both AdS and DL.

Observables

AdS/QCD

High Energy Physics – Theory [arXiv:1809.10536](#)

[Submitted on 27 Sep 2018]

Deep Learning and Holographic QCD

Koji Hashimoto, Sotaro Sugishita, Akinori Tanaka, Akio Tomiya

Phys. Rev. D **98**, 106014 – Published 14 November 2018

Chiral condensate

High Energy Physics – Theory [arXiv:2108.08091](#)

[Submitted on 18 Aug 2021 (v1), last revised 31 Aug 2021 (this version, v2)]

Deriving dilaton potential in improved holographic QCD from meson spectrum

Koji Hashimoto, Keisuke Ohashi, Takayuki Sumimoto

Meson spectrum

Phys. Rev. D **105**, 106008 – Published 13 May 2022

High Energy Physics – Phenomenology [arXiv:2401.06417](#)

[Submitted on 12 Jan 2024 (v1), last revised 30 Mar 2024 (this version, v2)]

Machine learning holographic black hole from lattice QCD equation of state

Xun Chen, Mei Huang

Lattice QCD

Phys. Rev. D **109**, L051902 – Published 25 March 2024

High Energy Physics – Theory [arXiv:2511.22522](#)

[Submitted on 27 Nov 2025]

AdS/Deep-Learning made easy II: neural network-based approaches to holography and inverse problems

Hyun-Sik Jeong, Hanse Kim, Keun-Young Kim, Gaya Yun, Hyeonwoo Yu, Kwan Yun

QCD Equations of State

AdS/CMT

High Energy Physics – Theory [arXiv:2004.12112](#)

[Submitted on 25 Apr 2020 (v1), last revised 3 Nov 2020 (this version, v3)]

Deep learning black hole metrics from shear viscosity

Yu-Kun Yan, Shao-Feng Wu, Xian-Hui Ge, Yu Tian

Phys. Rev. D **102**, 101902(R) – Published 12 November 2020

Shear viscosity

High Energy Physics – Theory [arXiv:2410.06523](#)

[Submitted on 9 Oct 2024]

Phase Diagram from Nonlinear Interaction between Superconducting Order and Density: Toward Data-Based Holographic Superconductor

Sejin Kim, Kyung Kiu Kim, Yunseok Seo

Superconductivity

J. High Energ. Phys. **2025**, 77 (2025)

High Energy Physics – Theory [arXiv:2401.00939](#)

[Submitted on 1 Jan 2024]

Deep learning bulk spacetime from boundary optical conductivity

Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

J. High Energ. Phys. **2024**, 141 (2024) **Optical conductivity**

High Energy Physics – Theory [arXiv:2502.10245](#)

[Submitted on 14 Feb 2025]

Deep learning-based holography for T-linear resistivity

Byoungjoon Ahn, Hyun-Sik Jeong, Chang-Woo Ji, Keun-Young Kim, Kwan Yun

Phys. Rev. D **112**, 126008 – Published 12 December, 2025

T-linear resistivity

High Energy Physics – Theory [arXiv:2411.16052](#)

[Submitted on 25 Nov 2024]

Machine-learning emergent spacetime from linear response in future tabletop quantum gravity experiments

Koji Hashimoto, Koshiro Matsuo, Masaki Murata, Gakuto Ogiwara, Daichi Takeda

[Machine Learning: Science and Technology](#), Volume 6, Number 1

AdS/QI

High Energy Physics – Theory [arXiv:2110.01115](#)

[Submitted on 3 Oct 2021]

Machine Learning Statistical Gravity from Multi-Region Entanglement Entropy

Jonathan Lam, Yi-Zhuang You

Phys. Rev. Research **3**, 043199 – Published 20 December 2021

High Energy Physics – Theory [arXiv:2205.04445](#)

[Submitted on 9 May 2022 (v1), last revised 11 Sep 2022 (this version, v3)]

Dual Geometry of Entanglement Entropy via Deep Learning

Chanyong Park, Chi-Ok Hwang, Kyungchan Cho, Se-Jin Kim

Phys. Rev. D **106**, 106017 – Published 28 November 2022

High Energy Physics – Theory [arXiv:2406.07395](#)

[Submitted on 11 Jun 2024 (v1), last revised 8 Sep 2024 (this version, v2)]

Holographic reconstruction of black hole spacetime: machine learning and entanglement entropy

Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

J. High Energ. Phys. **2025**, 25 (2025) **Entanglement entropy**

High Energy Physics – Theory [arXiv:2509.25311](#)

[Submitted on 29 Sep 2025]

Aspects of holographic entanglement using physics-informed-neural-networks

Anirudh Deb, Yaman Sanghavi

Wedge Cross Section

High Energy Physics – Theory [arXiv:2509.10866](#)

[Submitted on 13 Sep 2025 (v1), last revised 16 Sep 2025 (this version, v2)]

Physics-informed neural network solves minimal surfaces in curved spacetime

Koji Hashimoto, Koichi Kyo, Masaki Murata, Gakuto Ogiwara, Norihiro Tanahashi

MANY MORE ...

Observables

AdS/QCD

High Energy Physics – Theory [arXiv:1809.10536](#)

[Submitted on 27 Sep 2018]

Deep Learning and Holographic QCD

Koji Hashimoto, Sotaro Sugishita, Akinori Tanaka, Akio Tomiya

Phys. Rev. D **98**, 106014 – Published 14 November 2018

Chiral condensate

High Energy Physics – Theory [arXiv:2108.08091](#)

[Submitted on 18 Aug 2021 (v1), last revised 31 Aug 2021 (this version, v2)]

Deriving dilaton potential in improved holographic QCD from meson spectrum

Koji Hashimoto, Keisuke Ohashi, Takayuki Sumimoto

Meson spectrum

Phys. Rev. D **105**, 106008 – Published 13 May 2022

High Energy Physics – Phenomenology [arXiv:2401.06417](#)

[Submitted on 12 Jan 2024 (v1), last revised 30 Mar 2024 (this version, v2)]

Machine learning holographic black hole from lattice QCD equation of state

Xun Chen, Mei Huang

Lattice QCD

Phys. Rev. D **109**, L051902 – Published 25 March 2024

High Energy Physics – Theory [arXiv:2511.22522](#)

[Submitted on 27 Nov 2025]

AdS/Deep-Learning made easy II: neural network-based approaches to holography and inverse problems

Hyun-Sik Jeong, Hanse Kim, Keun-Young Kim, Gaya Yun, Hyeonwoo Yu, Kwan Yun

QCD Equations of State

AdS/CMT

High Energy Physics – Theory [arXiv:2004.12112](#)

[Submitted on 25 Apr 2020 (v1), last revised 3 Nov 2020 (this version, v3)]

Deep learning black hole metrics from shear viscosity

Yu-Kun Yan, Shao-Feng Wu, Xian-Hui Ge, Yu Tian

Phys. Rev. D **102**, 101902(R) – Published 12 November 2020

Shear viscosity

High Energy Physics – Theory [arXiv:2410.06523](#)

[Submitted on 9 Oct 2024]

Phase Diagram from Nonlinear Interaction between Superconducting Order and Density: Toward Data-Based Holographic Superconductor

Sejin Kim, Kyung Kiu Kim, Yunseok Seo

Superconductivity

J. High Energ. Phys. **2025**, 77 (2025)

High Energy Physics – Theory [arXiv:2401.00939](#)

[Submitted on 1 Jan 2024]

Deep learning bulk spacetime from boundary optical conductivity

Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

J. High Energ. Phys. **2024**, 141 (2024) **Optical conductivity**

High Energy Physics – Theory [arXiv:2502.10245](#)

[Submitted on 14 Feb 2025]

Deep learning-based holography for T-linear resistivity

Byoungjoon Ahn, Hyun-Sik Jeong, Chang-Woo Ji, Keun-Young Kim, Kwan Yun

Phys. Rev. D **112**, 126008 – Published 12 December, 2025

T-linear resistivity

High Energy Physics – Theory [arXiv:2411.16052](#)

[Submitted on 25 Nov 2024]

Machine-learning emergent spacetime from linear response in future tabletop quantum gravity experiments

Koji Hashimoto, Koshiro Matsuo, Masaki Murata, Gakuto Ogiwara, Daichi Takeda

[Machine Learning: Science and Technology](#), Volume 6, Number 1

AdS/QI

High Energy Physics – Theory [arXiv:2110.01115](#)

[Submitted on 3 Oct 2021]

Machine Learning Statistical Gravity from Multi-Region Entanglement Entropy

Jonathan Lam, Yi-Zhuang You

Phys. Rev. Research **3**, 043199 – Published 20 December 2021

High Energy Physics – Theory [arXiv:2205.04445](#)

[Submitted on 9 May 2022 (v1), last revised 11 Sep 2022 (this version, v3)]

Dual Geometry of Entanglement Entropy via Deep Learning

Chanyong Park, Chi-Ok Hwang, Kyungchan Cho, Se-Jin Kim

Phys. Rev. D **106**, 106017 – Published 28 November 2022

High Energy Physics – Theory [arXiv:2406.07395](#)

[Submitted on 11 Jun 2024 (v1), last revised 8 Sep 2024 (this version, v2)]

Holographic reconstruction of black hole spacetime: machine learning and entanglement entropy

Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

J. High Energ. Phys. **2025**, 25 (2025) **Entanglement entropy**

High Energy Physics – Theory [arXiv:2509.25311](#)

[Submitted on 29 Sep 2025]

Aspects of holographic entanglement using physics-informed-neural-networks

Anirudh Deb, Yaman Sanghavi

Wedge Cross Section

High Energy Physics – Theory [arXiv:2509.10866](#)

[Submitted on 13 Sep 2025 (v1), last revised 16 Sep 2025 (this version, v2)]

Physics-informed neural network solves minimal surfaces in curved spacetime

Koji Hashimoto, Koichi Kyo, Masaki Murata, Gakuto Ogiwara, Norihiro Tanahashi

MANY MORE ...

My Personal Goals (**Almost Dream, lol**)

Methodologies

Consistency

Can ML be a consistent methods?
(complement with traditional methods)

New Solutions

Can it find new solutions
not transparent in traditional methods?

Data

Robustness

How to construct models
that are both robust and accurate for data?

Predictions

What new insights and predictions
could ML provide?

- **OUTLINE**

1 | Methodology

: Physics-Informed Neural Networks (PINNs)

2 | Handful Examples

: AdS/QCD, CMT, QI

3 | Conclusion

Methodology


: Physics-Informed Neural Networks (PINNs)

Neural Networks

Nobel Prize in Physics 2024

“For foundational discoveries and inventions that enable machine learning with artificial neural networks”

Neural Networks




Training
parameters




$$y = \text{NN}(x, \theta)$$

Neural networks are simply **flexible functions** fit to data




Given data, tune the **parameter** θ so that the network approximates the true function

$$\text{NN}(x, \theta) \approx y_{\text{true}}(x)$$



Biological Neuron


Perceptron





Training parameters
↓

$$y = \text{NN}(x, \theta)$$

Perceptron is the fundamental unit of a neural network



Activation Function




$$\text{IF, } \sum_i w_i x_i \geq \theta \longrightarrow y = 1$$

(Threshold) (Activated!)


$$\text{IF, } \sum_i w_i x_i < \theta \longrightarrow y = 0$$

(Deactivated!)

Activation function adds non-linearity by changing the output of a neuron, helping the model capture more complex patterns



Deep Neural Networks



Deep neural networks are composition functions of NNs

$$\text{NN}(x, \theta) \approx W_M \cdot f[\cdots f\{\underbrace{W_2 \cdot f(W_1 x + b_1)}_x + b_2\} \cdots] + b_M$$

$\theta = \{W_i, b_i\}$ are learning parameters

of layers

Universal Approximation Theorem

Original contribution Neural Networks Volume 2, Issue 5, 1989, Pages 359-366

Multilayer feedforward networks are universal approximators

Kurt Hornik, Maxwell Stinchcombe, Halbert White¹ 



This paper rigorously establishes that standard multilayer feedforward networks with as few as one hidden layer using arbitrary squashing functions are capable of approximating any Borel measurable function from one finite dimensional space to another to any desired degree of accuracy, provided sufficiently many hidden units are available. In this sense, multilayer feedforward networks are a class of universal approximators.

DNNs, $\text{NN}(x, \theta)$, can approximate
any continuous functions $y(x)$



How to find such solutions? or how to train DNNs?

Existence Theorem

Definition 2.1


For any $r \in \mathbb{N} \equiv \{1, 2, \dots\}$, A^r is the set of all affine functions from $\mathbb{R}^r$ to $\mathbb{R}$, that is, the set of all functions of the form $A(x) = w \cdot x + b$ where w and x are vectors in $\mathbb{R}^r$, “ $\cdot$ ” denotes the usual dot product of vectors, and $b \in \mathbb{R}$ is a scalar. $\square$



For every function g in M^r there is a compact subset K of $\mathbb{R}^r$ and an $f \in \Sigma^r(\Psi)$ such that for any $\varepsilon > 0$ we have $\mu(K) < 1 - \varepsilon$ and for every $x \in K$ we have $|f(x) - g(x)| < \varepsilon$, regardless of Ψ , r , or μ . $\square$


$$|y(x) - \text{NN}(x, \theta)| < \epsilon$$

Loss Function



We tune the parameters **minimizing** some loss function $L(\theta)$

Example:
Mean Squared Error
(L2 Loss)

$$L(\theta) = \frac{1}{N} \sum_i^N [\text{NN}(x_i, \theta) - y(x_i)]^2$$

Training DNNs


Data

Typically, by using **gradient decent**,
parameters are optimized iteratively


$$\theta^{j+1} = \theta^j - l_r \nabla_{\theta} L(\theta^j)$$

the iteration (epoch)

learning rate (step size)



Loss Function



We tune the parameters **minimizing** some loss function $L(\theta)$

Example:
Mean Squared Error
(L2 Loss)

$$L(\theta) = \frac{1}{N} \sum_i^N [\text{NN}(x_i, \theta) - y(x_i)]^2$$

Training DNNs


Data

Typically, by using **gradient decent**,
parameters are optimized iteratively

$$\theta^{j+1} = \theta^j - l_r \nabla_{\theta} L(\theta^j)$$


the iteration (epoch)

learning rate (step size)





Deep Neural Networks
Automatic Differentiation
Adam Optimizer L-BFGS



We tune the parameters **minimizing** some loss function $L(\theta)$

Example:
Mean Squared Error
(L2 Loss)

$$L(\theta) = \frac{1}{N} \sum_i^N [\text{NN}(x_i, \theta) - y(x_i)]^2$$


Training DNNs Data

Typically, by using **gradient decent**,
parameters are optimized iteratively

$$\theta^{j+1} = \theta^j - l_r \nabla_{\theta} L(\theta^j)$$

the iteration (epoch)

learning rate (step size)



Physics-Informed Neural Networks

(PINNs)

Physics-Informed Neural Networks

Journal of Computational Physics Volume 378, 1 February 2019, Pages 686-707

Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations

M. Raissi^a, P. Perdikaris^b , G.E. Karniadakis^a



CITED BY YEAR

18453

2019

PINNs incorporate **physical laws (PDE, ODEs)**, directly into the training process.

$$L(\theta) = L(\theta)_{\text{Data}} + L(\theta)_{\text{Equations}}$$

Standard NN
(solely rely on data)

Data Loss

To ensure the model
fits the available data points




Physical Loss

To encode the governing
physical equations

- 1) to construct **robust models** (even with limited or noisy data) preventing overfitting problem
- 2) a **new (forward problem) solver** to model system dynamics
- 3) an **inverse problem solver** to learn the underlying physical parameters of the system

Data
+
Eqns



Broad Applicability of PINNs

[Home](#) > [Journal of Scientific Computing](#) > [Article](#)

Scientific Machine Learning Through Physics-Informed Neural Networks: Where we are and What's Next

[Open access](#) | Published: 26 July 2022

Volume 92, article number 88, (2022) [Cite this article](#)

Computer Science > Machine Learning

arXiv:2410.13228

From PINNs to PIKANs: Recent Advances in Physics-Informed Machine Learning

Juan Diego Toscano^{a,1}, Vivek Oommen^{b,1}, Alan John Varghese^{b,1}, Zongren Zou^a,
Nazanin Ahmadi Daryakenari^{b,c}, Chenxi Wu^b, George Em Karniadakis^{a,b,2}

^aDivision of Applied Mathematics, Brown University, Providence, 02912, RI, USA

^bSchool of Engineering, Brown University, Providence, 02912, RI, USA

^cCenter for Biomedical Engineering, Brown University, Providence, 02912, RI, USA

Computer Science > Machine Learning

arXiv:2510.05433

Physics-Informed Machine Learning in Biomedical Science and Engineering

[Home](#) > [Acta Mechanica Sinica](#) > [Article](#)

Physics-informed neural networks (PINNs) for fluid mechanics: a review

Invited Review | Published: 23 January 2022

[Journals & Magazines](#) > [IEEE Transactions on Power Sy...](#) > Volume: 38 Issue: 1

Applications of Physics-Informed Neural Networks in Power Systems - A Review

Publisher: IEEE

[Cite This](#)

[PDF](#)


Bin Huang ; Jianhui Wang [All Authors](#)

Hidden fluid mechanics: Learning velocity and pressure fields from flow visualizations

MAZIAR RAISSI , ALIREZA YAZDANI , AND GEORGE EM KARNIADAKIS  [Authors Info & Affiliations](#)

SCIENCE • 30 Jan 2020 • Vol 367, Issue 6481 • pp. 1026-1030 • DOI: 10.1126/science.aaw4741

**Data +
Navier-Stokes Eqns**




Example

Damped Harmonic Oscillator


$$x''(t) + 2\delta x'(t) + \omega_0^2 x(t) = 0$$

Damping coefficient

Spring constant




Data Preparation



PyTorch

```
import torch
import torch.nn as nn
```

nn: Neural Networks



DNNs

```
def __init__(self, N_INPUT, N_OUTPUT, N_HIDDEN, N_LAYERS):
    super().__init__()
    activation = nn.Tanh
    self.fcs = nn.Sequential(*[nn.Linear(N_INPUT, N_HIDDEN), activation()])
    self.fch = nn.Sequential(*[nn.Sequential(*[nn.Linear(N_HIDDEN, N_HIDDEN), activation()]) for _ in range(N_LAYERS-1)])
    self.fce = nn.Linear(N_HIDDEN, N_OUTPUT)
```

δ
 ω_0

```
d = torch.nn.Parameter(torch.zeros(1, requires_grad=True))
w0 = torch.nn.Parameter(torch.empty(1).uniform_(0, 50), requires_grad=True)
```

```
torch.optim.Adam(list(pinn.parameters())+[d]+[w0], lr=1e-3)
```

```
dxdt = torch.autograd.grad(x, t_physics, torch.ones_like(x), create_graph=True)[0]
```

Training
parameters

Optimizer

Auto-grad


Loss

$$\mathcal{L}(\theta, \delta, \omega_0) = \frac{1}{N} \sum_i^N [x''_{\text{PINN}}(t_i; \theta) + 2\delta x'_{\text{PINN}}(t_i; \theta) + \omega_0^2 x_{\text{PINN}}(t_i; \theta)]^2 + \frac{\lambda}{M} \sum_j^M (x_{\text{PINN}}(t_j; \theta) - x_{\text{obs}}(t_j))^2$$

Physical Loss

Data Loss

1) to construct **robust models** (even with limited or noisy data) preventing
overfitting problem




Standard Neural Network solution

(training only with **data**)

“It’s suffering from the overfitting problem even in simple example”

“It’s quite poor outside the data domain”



Physics-Informed Neural Network solution

(training with **data + equations**)

“PINN allows accurate predictions even with limited training data”


“Working well even outside the data domain”

2) a **new (forward problem) solver** to model system dynamics


$$\mathcal{L}(\theta) = \underbrace{[x_{\text{PINN}}(t=0; \theta) - 1]^2 + \lambda_1 [x'_{\text{PINN}}(t=0; \theta) - 0]^2}_{\text{Boundary Conditions}} + \underbrace{\frac{\lambda_2}{N} \sum_i^N [x''_{\text{PINN}}(t_i; \theta) + 2\delta x'_{\text{PINN}}(t_i; \theta) + \omega_0^2 x_{\text{PINN}}(t_i; \theta)]^2}_{\text{Equation of Motion}}$$

Boundary Conditions

Equation of Motion



Training
→
Minimizing
Loss
Function



$$x_{\text{Exact}}(t) = e^{-\delta t} (c_1 e^{i\omega t} + c_2 e^{-i\omega t})$$

$$(\delta, \omega_0) = (1, 20)$$


$$\omega := \sqrt{\omega_0^2 - \delta^2}$$

3) an **inverse problem solver** to learn the underlying physical parameters of the system




$$\mathcal{L}(\theta, \delta, \omega_0) = \underbrace{\frac{\lambda}{M} \sum_j^M (x_{\text{PINN}}(t_j; \theta) - x_{\text{obs}}(t_j))^2}_{\text{Data or B.C.s}} + \underbrace{\frac{1}{N} \sum_i^N [x_{\text{PINN}}''(t_i; \theta) + 2\delta x_{\text{PINN}}'(t_i; \theta) + \omega_0^2 x_{\text{PINN}}(t_i; \theta)]^2}_{\text{Equation of Motion}}$$

Data or B.C.s

Equation of Motion



Training



3) an **inverse problem solver** to learn the underlying physical parameters of the system



$$\mathcal{L}(\theta, \delta, \omega_0) = \underbrace{\frac{\lambda}{M} \sum_j^M (x_{\text{PINN}}(t_j; \theta) - x_{\text{obs}}(t_j))^2}_{\text{Data or B.C.s}} + \underbrace{\frac{1}{N} \sum_i^N [x''_{\text{PINN}}(t_i; \theta) + 2\delta x'_{\text{PINN}}(t_i; \theta) + \omega_0^2 V(x)]^2}_{\text{Equation of Motion}}$$

Data or B.C.s



Equation of Motion

Finding potentials !!

Training step 15000



Estimated $V(x)$ vs True $V(x)$



Handful Examples

AdS/QCD, CMT, QI

AdS/QCD, CMT

arXiv:2401.00939 [pdf, other] hep-th cond-mat.dis-nn gr-qc hep-ph doi 10.1007/JHEP03(2024)141

Deep learning bulk spacetime from boundary optical conductivity

Authors: Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

arXiv:2502.10245 [pdf, other] hep-th cond-mat.dis-nn cond-mat.str-el gr-qc doi 10.1103/nh4n-ylnz (PRD)

Deep learning-based holography for T-linear resistivity

Authors: Byoungjoon Ahn, Hyun-Sik Jeong, Chang-Woo Ji, Keun-Young Kim, Kwan Yun

arXiv:2511.22522 [pdf, ps, other] hep-th cs.LG

AdS/Deep-Learning made easy II: neural network-based approaches to holography and inverse problems

Authors: Hyun-Sik Jeong, Hanse Kim, Keun-Young Kim, Gaya Yun, Hyeonwoo Yu, Kwan Yun

Setup

Action

$$S = \int d^4x \sqrt{-g} (R + \mathcal{L}_m) , \quad \mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + \boxed{V(\phi)} - \boxed{\frac{Z(\phi)}{4}} F^2 - \frac{1}{2} \sum_{i=1}^2 (\partial\chi_i)^2$$


**Bulk
Fields**

$$ds^2 = \frac{1}{r^2} \left(\boxed{-f(r)} dt^2 + \frac{1}{f(r)} dr^2 + \boxed{h(r)} \sum_{i=1}^2 dx_i^2 \right)$$

$$\phi = \boxed{\phi(r)}, \quad A = \boxed{A_t(r)} dt, \quad \chi_i = \beta x_i,$$

“to break translational sym.”

DNNs



Analogy

$$x(t) \rightarrow \boxed{g_{\mu\nu}(r), A_t(r), \phi(r)}$$

$$V(x) \rightarrow \boxed{V(\phi), Z(\phi)}$$

$$\{\delta, \omega_0\} \rightarrow \beta$$

Damped Harmonic Oscillator

$$x''(t) + 2\delta x'(t) + \omega_0^2 V(x) = 0$$

Setup

$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Loss Function


$$L = L_{\text{Data}} + L_{\text{EOM}}$$

Einstein's equations,
Maxwell' equations,
Dilaton equations

QCD Equation of State (Speed of Sound)

“Black hole thermodynamics”

$$C_S^2 = \frac{d \log T}{d \log s}$$




“we give horizon constraints”

Optical Conductivity

“Two-point functions”

$$\sigma(\omega) = \frac{1}{i\omega} G_{j^x j^x}^R(\omega) = \frac{1}{i\omega} \frac{a_x^{(R)}}{a_x^{(S)}}$$




$$a_x = a_x^{(S)} + a_x^{(R)} z + \dots$$

T-linear Resistivity

$$\rho = \frac{1}{\sigma_{DC}}$$

$$\sigma_{DC} = Z_H + \frac{q^2}{\beta^2 h(1)}$$

$$q = Z_H h(1) A'_t(1)$$



“we give horizon constraints”

Results

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.

I. QCD Equation of State (Speed of Sound)

PRL **101**, 131601 (2008)

PHYSICAL REVIEW LETTERS

week ending
26 SEPTEMBER 2008

Thermodynamics and Bulk Viscosity of Approximate Black Hole Duals to Finite Temperature Quantum Chromodynamics

Steven S. Gubser,^{*} Abhinav Nellore,⁺ Silviu S. Pufu,[‡] and Fábio D. Rocha[§]
Joseph Henry Laboratories, Princeton University, Princeton, New Jersey 08544, USA
(Received 30 June 2008; published 23 September 2008)

PHYSICAL REVIEW D **78**, 086007 (2008)

Mimicking the QCD equation of state with a dual black hole


Steven S. Gubser and Abhinav Nellore
Joseph Henry Laboratories, Princeton University, Princeton, New Jersey 08544, USA
(Received 25 June 2008; published 29 October 2008)

$$\int d^5x \sqrt{-g} \left(R - \frac{1}{2} (\partial \phi)^2 - V(\phi) \right)$$

“By human learning (trial-error)”


$$V(\Phi) = -12 \cosh(\gamma \Phi) + b \Phi^2,$$

$$\text{with } \gamma = 0.606, \quad b = 2.057$$



Results

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.







$$\int d^5x \sqrt{-g} \left(R - \frac{1}{2} (\partial \phi)^2 - V(\phi) \right)$$

“By human learning (trial-error)”

$$V(\Phi) = -12 \cosh(\gamma \Phi) + b \Phi^2,$$

with $\gamma = 0.606$, $b = 2.057$



Results

6 1


$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.


II. Optical Conductivity

$$\int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4}F_{ab}F^{ab} - \frac{1}{2}\sum_{I=1}^2(\partial X_I)^2 \right)$$

“Linear-axion model”: the effective toy model



$$\chi_i = \beta x_i$$



Holographic Optical Conductivity

[Home](#) > [Journal of High Energy Physics](#) > [Article](#)

A simple holographic model of momentum relaxation

[Open access](#) | Published: 22 May 2014

Volume 2014, article number 101, (2014) [Cite this article](#)

[Tomás Andrade](#) ✉ & [Benjamin Withers](#)

[Home](#) > [Science China Physics, Mechanics & Astronomy](#) > [Article](#)

Holographic axion model: A simple gravitational tool for quantum matter

Invited Review | Published: 01 June 2021

Volume 64, article number 270001, (2021) [Cite this article](#)

[Matteo Baggioli](#), [Keun-Young Kim](#) ✉, [Li Li](#) ✉ & [Wei-Jia Li](#) ✉


Results

$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.

II. Optical Conductivity

“Emblackening Factor”




■ Dots = True Solutions

— Lines = PINNs


$$\int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4}F_{ab}F^{ab} - \frac{1}{2}\sum_{I=1}^2(\partial X_I)^2 \right)$$

“Linear-axion model”: the effective toy model



$$\chi_i = \beta x_i$$

Holographic Optical Conductivity



Results

$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.

II. Optical Conductivity


nature

Letter | Published: 22 December 2005


Extremely slow Drude relaxation of correlated electrons

Marc Scheffler , Martin Dressel, Martin Jourdan & Hermann Adrian

Nature **438**, 1135–1137 (2005) | [Cite this article](#)



“Heavy Fermion Metals”



Linear-axion
model with

$$\beta = 0.4825$$

$$\sigma(\omega) = \frac{\sigma_0}{1 - i\omega\tau}$$

```
δ = torch.nn.Parameter(torch.zeros(1, requires_grad=True))
ω₀ = torch.nn.Parameter(torch.empty(1).uniform_(0, 50), requires_grad=True)
```


Results

$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.

III. T-linear Resistivity

PHYSICAL REVIEW D **81**, 046001 (2010)

Peculiar properties of a charged dilatonic black hole in AdS₅

Steven S. Gubser and Fabio D. Rocha

Joseph Henry Laboratories, Princeton University, Princeton, New Jersey 08544, USA
(Received 30 November 2009; published 1 February 2010)

PHYSICAL REVIEW B **89**, 245116 (2014)

Holographic duality and the resistivity of strange metals

Richard A. Davison,^{1,*} Koenraad Schalm,^{1,2,†} and Jan Zaanen^{1,‡}

¹*Instituut-Lorentz for Theoretical Physics, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands*






²*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

(Received 12 November 2013; revised manuscript received 17 May 2014; published 12 June 2014)




$$V(\phi) = 6 \cosh \frac{\phi}{\sqrt{3}}, \quad Z(\phi) = e^{\frac{\phi}{\sqrt{3}}},$$

“Gubser-Rocha model”

- 1) Top-down string theory models
- 2) It allows the analytic background solutions
- 3) It shows T-linear resistivity and linear specific heat
(as in cuprate strange metals)




Training



Results


$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.



$$V(\phi) \approx e^{\frac{1}{\sqrt{3}}\phi}$$


Training



$$Z(\phi) \approx \alpha e^{\frac{1}{\sqrt{3}}\phi}$$

$$\alpha = \alpha(\mathcal{A})$$

Gubser-Rocha model



$$\rho = \mathcal{A} T$$

Results

$$\mathcal{L}_m = -\frac{1}{2}(\partial\phi)^2 + V(\phi) - \frac{Z(\phi)}{4}F^2 - \frac{1}{2}\sum_{i=1}^2(\partial\chi_i)^2$$

Mostly, we haven't found anything new yet, but **the consistency** with standard holography results, and **some interesting observations**.

yses. Specifically, one can check that within the EMDA theories (3.1) [24–27, 36–43], where the axion-dilaton coupling $\frac{Y(\phi)}{2}\sum_{i=1}^2(\partial\chi_i)^2$ is simplified to $Y(\phi) = 1$, the corresponding IR geometry takes the form of the called η geometry:

$$ds^2 = \xi^{-\eta} \left(\frac{-dt^2 + \mathcal{L}^2 d\xi^2}{\xi^2} + \sum_{i=1}^2 dx_i^2 \right), \quad \mathcal{L}^2 := \frac{2(1+\eta)^2}{2-\beta^2}. \quad (4.1)$$

Here, the scaling parameters satisfy

$$z \rightarrow \infty, \quad \theta \rightarrow -\infty, \quad \theta/z \rightarrow -\eta. \quad V_{\text{IR}}(\phi) \approx Z_{\text{IR}}(\phi) \approx e^{\frac{\phi}{\sqrt{3}}}$$

In this geometry, the resistivity and entropy scale at low T as

$$\rho \approx T^\eta, \quad s \approx T^\eta.$$

**Consistent with the
IR Geometry Analysis**

AdS/QI

[arXiv:2406.07395](#) [pdf, other] hep-th cond-mat.dis-nn gr-qc doi [10.1007/JHEP01\(2025\)025](#)


Holographic reconstruction of black hole spacetime: machine learning and entanglement entropy

Authors: Byoungjoon Ahn, Hyun-Sik Jeong, Keun-Young Kim, Kwan Yun

Setup

$$S = \frac{\text{Area}(\gamma)}{4G_N}$$

Bulk Reconstruction



Action

We do not need the action

**Bulk
Fields**

$$ds^2 = \frac{L^2}{z^2} \left[-f(z) dt^2 + \frac{dz^2}{f(z)} + h(z) (dx^2 + dy^2) \right]$$

“General homogenous (isotropic) metric”


“Coordinate transformation”

$$ds^2 = -D(r) dt^2 + B(r) dr^2 + C(r) d\vec{x}_i^2$$

DNNs



= z





f or h

Holographic Entanglement Entropy

Entanglement entropy in the boundary theory can be computed from **the extremal surface** in the higher dimensional gravity theory

$$S = \frac{\text{Area}(\gamma)}{4G_N}$$
$$= S(\ell)$$



Holographic Entanglement Entropy

Entanglement entropy in the boundary theory can be computed from **the extremal surface** in the higher dimensional gravity theory

$$S = \frac{\text{Area}(\gamma)}{4G_N}$$

$$= S(\ell)$$

$$\text{Area}(\gamma) = \int dy dx \sqrt{\det \gamma_{\mu\nu}}$$

“Induced metric at time slice”

“Finding the ext. area by Euler-Lagrangian method”

$$ds^2 = \frac{L^2}{z^2} \left[-f(z) dt^2 + \frac{dz^2}{f(z)} + h(z) (dx^2 + dy^2) \right]$$


$$S = \frac{L^2 \Omega}{2G_N} \int_{\epsilon}^{z_*} \frac{1}{z^2} \sqrt{\frac{h(z)}{1 - \frac{z_*^4 h(z_*)^2}{z^4 h(z)^2}}} \frac{1}{\sqrt{f(z)}} dz,$$

$$\ell = 2 \int_0^{z_*} \frac{1}{\sqrt{\frac{h(\alpha)^2}{h(z_*)^2} \frac{z_*^4}{\alpha^4} - 1}} \frac{1}{\sqrt{h(\alpha) f(\alpha)}} d\alpha,$$

One more thing...

Example: 1+1 CFT $S_A = \frac{c}{3} \log \frac{\ell}{\epsilon} + \text{finite}$

central charge of the CFT
UV cut off



$$S = \frac{L^2 \Omega}{2G_N} \int_{\epsilon}^{z_*} \frac{1}{z^2} \sqrt{\frac{h(z)}{1 - \frac{z^4 h(z_*)^2}{z_*^4 h(z)^2}}} \frac{1}{\sqrt{f(z)}} dz$$


$$= \frac{L^2 \Omega}{2G_N} \left[\int_{\epsilon}^{z_*} \frac{1}{z^2} dz + \int_0^{z_*} \frac{1}{z^2} \left(\sqrt{\frac{h(z)}{1 - \frac{z^4 h(z_*)^2}{z_*^4 h(z)^2}}} \frac{1}{\sqrt{f(z)}} - 1 \right) dz \right]$$

$$= \frac{L^2 \Omega}{2G_N} \left[\frac{1}{\epsilon} + \left\{ \int_0^{z_*} \frac{1}{z^2} \left(\sqrt{\frac{h(z)}{1 - \frac{z^4 h(z_*)^2}{z_*^4 h(z)^2}}} \frac{1}{\sqrt{f(z)}} - 1 \right) dz - \frac{1}{z_*} \right\} \right]$$

$$:= \frac{L^2 \Omega}{2G_N} \left[\frac{1}{\epsilon} + S_{\text{Finite}} \right],$$

**Loss
Function**

$$L = \frac{1}{N_{\ell}} \sum_{\ell} \left| S_{\text{Finite}}^{(m)}(\ell) - S_{\text{Finite}}(\ell) \right| + \left| s^{(m)} - s \right|$$



Technical Reason
(Must satisfy by construction)

Bulk Reconstruction

(Without Machine Learning)

$$S = \frac{L^2 \Omega}{2G_N} \int_{\epsilon}^{z_*} \frac{1}{z^2} \sqrt{\frac{h(z)}{1 - \frac{z^4}{z_*^4} \frac{h(z_*)^2}{h(z)^2}}} \frac{1}{\sqrt{f(z)}} dz$$

Handbook of
integral eqns

Output

$$F(z_*) = \int_{\epsilon}^{z_*} \frac{1}{\sqrt{G(z_*) - G(z)}} Y(z) dz, \quad G'(z_*) > 0$$

Input

(Solution) :

$$Y(z_*) = \frac{1}{\pi} \frac{d}{dz_*} \int_{\epsilon}^{z_*} \frac{G'(z)}{\sqrt{G(z_*) - G(z)}} F(z) dz.$$

For the simple
metric

$$h(z) = 1$$

$$ds^2 = \frac{L^2}{z^2} \left[-f(z) dt^2 + \frac{dz^2}{f(z)} + h(z) (dx^2 + dy^2) \right]$$

$$\frac{2G_N}{L^2 \Omega} \frac{S(z_*)}{z_*^2} = \int_{\epsilon}^{z_*} \frac{1}{\sqrt{z_*^4 - z^4}} \frac{\sqrt{1/f(z)}}{z^2} dz$$

$$G(z) := z^4$$

$$\frac{dS(\ell)}{d\ell} = \frac{L^2 \Omega}{4G_N} \frac{1}{z_*^2}$$

$$\sqrt{\frac{1}{f(z)}} = \frac{8G_N}{\pi L^2 \Omega} z^2 \frac{d}{dz} \int_{\epsilon}^z \frac{z_*}{\sqrt{z^4 - z_*^4}} S(z_*) dz_*$$

We **do not need ML** to reconstruct the bulk from EE **when metric is simple**


Bulk Reconstruction

(Example)

$$h(z) = 1$$

$$f(z) = 1 - \frac{\beta^2}{2} z^2 - \left(1 - \frac{\beta^2}{2} + \frac{\mu^2}{4}\right) z^3 + \frac{\mu^2}{4} z^4$$

Given holographic
entanglement entropy




$$\int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F_{ab} F^{ab} - \frac{1}{2} \sum_{I=1}^2 (\partial X_I)^2 \right)$$

“Linear-axion model”: the effective toy model

$$\sqrt{\frac{1}{f(z)}} = \frac{8G_N}{\pi L^2 \Omega} z^2 \frac{d}{dz} \int_{\epsilon}^z \frac{z_*}{\sqrt{z^4 - z_*^4}} S(z_*) dz_*$$

Bulk reconstruction
without machine learning



Bulk Reconstruction

(Limitation)

$$S = \frac{L^2 \Omega}{2G_N} \int_{\epsilon}^{z_*} \frac{1}{z^2} \sqrt{\frac{h(z)}{1 - \frac{z^4 h(z_*)^2}{z_*^4 h(z)^2}}} \frac{1}{\sqrt{f(z)}} dz$$

For the **generic** metric, **ML becomes very powerful** (necessary) for reconstruction

$h(z) \neq 1$ (This is very very common situation in holography)

$$ds^2 = \frac{L^2}{z^2} \left[-f(z) dt^2 + \frac{dz^2}{f(z)} + h(z) (dx^2 + dy^2) \right]$$

Same
Information

$S(z_*)$

(See details in our paper)

Output

$$\sqrt{\frac{1}{f(z) h(z)^3}} = \frac{2}{\pi} \frac{1}{z^2} \frac{d}{dz} \int_0^z \frac{\ell(z_*)}{\sqrt{\frac{z^4}{h(z)^2} - \frac{z_*^4}{h(z_*)^2}}} \frac{2z_*^3 h(z_*) - z_*^4 h'(z_*)}{2h(z_*)^3} dz_*$$

Single information



But, two outputs

$$\frac{dS}{d\ell} = \frac{L^2 \Omega}{4G_N} \frac{h(z_*)}{z_*^2}$$

“Unless $h(z)$ is provided, this method cannot fully reconstruct the metric”


Bulk Reconstruction

(Machine Learning Example I)

$$\int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F_{ab} F^{ab} - \frac{1}{2} \sum_{I=1}^2 (\partial X_I)^2 \right)$$

“Linear-axion model”: the effective toy model



Given holographic
entanglement entropy



■ Dots = True Solutions

— Lines = PINNs


Bulk reconstruction
with machine learning



Bulk Reconstruction

(Machine Learning Example II)

Given holographic
entanglement entropy





■ Dots = True Solutions

— Lines = PINNs

$$V(\phi) = 6 \cosh \frac{\phi}{\sqrt{3}}, \quad Z(\phi) = e^{\frac{\phi}{\sqrt{3}}},$$

“Gubser-Rocha model”

Bulk reconstruction
with machine learning



$h(z) \neq 1$



Bulk Reconstruction

(Machine Learning Example III)



$$S = \int d^4x \sqrt{-g} \left(R + 6 - \frac{1}{4} F^2 - |D\Phi|^2 - m^2 |\Phi|^2 \right)$$

“Holographic Superconductor”

Given holographic
entanglement entropy



Bulk reconstruction
with machine learning



■ Dots = True Solutions

— Lines = PINNs

Bulk Reconstruction

(**Machine Learning** Example IV)

$$ds^2 = \frac{L^2}{z^2} \left[-f(z) dt^2 + \frac{dz^2}{f(z)} + h(z) dx^2 \right]$$


“AdS₃ metric” $\longleftrightarrow$ “(1+1) dimensional system”

$$S = \frac{L}{2G_N} \log \left[\frac{2}{\epsilon} \sinh \left(\frac{\ell}{2} \right) \right]$$

$$S_{\text{CFT}} = \frac{c}{3} \log \left[\frac{\beta}{\pi \epsilon} \sinh \left(\frac{\pi \ell}{\beta} \right) \right]$$

Given **non-holographic**
entanglement entropy

??



Bulk reconstruction
with machine learning

PHYSICAL REVIEW LETTERS 127, 040603 (2021)

Eigenstate Entanglement: Crossover from the Ground State to Volume Laws

Qiang Miao and Thomas Barthel

Department of Physics, Duke University, Durham, North Carolina 27708, USA

(Received 30 May 2019; revised 27 May 2021; accepted 14 June 2021; published 21 July 2021)

To confirm this numerically, we sample energy eigenstates $|E_n\rangle$ from small windows of width ΔE around energies E for fermionic tight-binding chains $\hat{H} = -\sum_i (\hat{c}_i^\dagger \hat{c}_{i+1} + \text{H.c.}) = \sum_k \epsilon_k \hat{n}_k$ at half filling [65]. These obey weak ETH and are

Similar to CFT₂ result
(so, similar to BTZ as well?)

$$\hat{\rho}_n := \text{Tr}_B |E_n\rangle \langle E_n|$$






FIG. 1. Entanglement entropies of randomly sampled eigenstates for a fermionic tight-binding chain with half filling ($\mu = 0$, $v = 2$), system size $L \approx 4 \times 10^6$, excitation energy densities $e := (E - E_{\text{gs}})/|E_{\text{gs}}|$, and window size $\Delta E = 1$. The main panel confirms the data collapse to the crossover scaling function (7). The inset asserts the validity of the ETH by comparison to GCE subsystem entropies (lines).

Bulk Reconstruction

(Machine Learning Example IV)





“Gubser-Rocha model”

Given **non-holographic**
entanglement entropy

??




Bulk reconstruction
with machine learning



The emergent bulk spacetime is not similar to BTZ geometries, instead..
Gubser-Rocha type.. the similarity is due to the metallic property??




Conclusion

Neural networks are simply **flexible functions** fit to data




Modeling Realistic Systems

Bulk Reconstruction





**Potential implications
are far-reaching !!!**



THANK YOU

Hyun-Sik Jeong

APCTP

apctp asia pacific center for
theoretical physics